International Journal of Modern Physics A
© World Scientific Publishing Company

arXiv:hep-ph/0509316v2 13 Oct 2005

HADRON TOMOGRAPHY

MATTHIAS BURKARDT
Department of Physics, New Mexico State University
Las Cruces, NM 88011, USA

Received (Day Month Year)
Revised (Day Month Year)

Generalized parton distributions can be used to obtain information about the dependence of parton distributions on the impact parameter. Potential consequences for T-odd single-spin asymmetries are discussed.

Keywords: GPDs; SSA

1. Introduction

Hadron form factors provide information about the Fourier transform of the charge distribution within the hadron. Generalized parton distributions (GPDs) provide a momentum decomposition of the form factor w.r.t. the average momentum fraction $x = \frac{1}{2}(x_i + x_f)$ of the active quark

$$\int dx H_q(x,\xi,t) = F_1^q(t) \qquad \int dx E_q(x,\xi,t) = F_2^q(t) \tag{1}$$

where $F_1^q(t)$ and $F_2^q(t)$are the Dirac and Pauli formfactors, respectively. x_i and x_f are the momentum fractions of the quark before and after the momentum transfer.

The momentum direction of the active quark singles out a direction in space and therefore it makes a difference whether the momentum transfer is along that momentum or in a different direction. Therefore GPDs not only depend on x and the invariant momentum transfer t but also on the longitudinal momentum transfer through the variable $2\xi = x_f - x_i$. Unlike form factors, in which the contribution from all quark momenta are always summed up, GPDs thus tell us how much each momentum contributes to the form factor at a given momentum transfer.

GPDs are the form factor of the same operator whose forward matrix elements yield the usual parton distribution functions (PDFs)

$$\int \frac{dx^-}{2\pi} e^{ix^-\bar{p}^+x} \left\langle p' \left| \bar{q}\left(-\frac{x^-}{2}\right)\gamma^+ q\left(\frac{x^-}{2}\right)\right| p\right\rangle = H(x,\xi,\Delta^2)\bar{u}(p')\gamma^+u(p) \tag{2}$$
$$+E(x,\xi,\Delta^2)\bar{u}(p')\frac{i\sigma^{+\nu}\Delta_\nu}{2M}u(p),$$

This observation has formed the basis of the position space interpretation of GPDs.

2. Position Space Interpretation of GPDs

Charge distributions in position space are usually measured in the center of mass frame, i.e. measured relative to the center of mass of the system. For impact parameter dependent parton distributions, the analogous reference point is the transverse center of momentum of all partons (quarks and gluons) $\mathbf{R}_\perp = \sum_{i=q,g} x_i \mathbf{r}_{\perp,i}$, where x_i is the momentum fraction carried by each parton and $\mathbf{r}_{\perp,i}$ is their $\perp$ position. This $\perp$ center of momentum behaves in many ways similar to the nonrelativistic center of mass. For example, one can form eigenstates of $\mathbf{R}_\perp$

$$\left|p^+, \mathbf{R}_\perp = \mathbf{0}_\perp, \lambda\right\rangle \equiv \mathcal{N} \int d^2\mathbf{p}_\perp \left|p^+, \mathbf{p}_\perp, \lambda\right\rangle . \tag{3}$$

Impact parameter dependent PDFs are defined using the familiar light-cone correlation function in such transversely localized states

$$q(x, \mathbf{b}_\perp) \equiv \int \frac{dx^-}{4\pi} \left\langle p^+, \mathbf{R}_\perp = \mathbf{0}_\perp\right| \bar{q}(-\frac{x^-}{2}, \mathbf{b}_\perp)\gamma^+ q(\frac{x^-}{2}, \mathbf{b}_\perp) \left|p^+, \mathbf{R}_\perp = \mathbf{0}_\perp\right\rangle e^{ixp^+x^-}, \tag{4}$$

and similarly for the polarized distribution $\Delta q(x, \mathbf{b}_\perp)$ with an additional γ_5.

Impact parameter dependent PDFs are Fourier transforms of GPDs for $\xi = 0$

$$q(x, \mathbf{b}_\perp) = \int \frac{d^2\mathbf{\Delta}_\perp}{(2\pi)^2} e^{i\mathbf{\Delta}_\perp \cdot \mathbf{b}_\perp} H(x, 0, -\mathbf{\Delta}_\perp^2) \tag{5}$$

$$\Delta q(x, \mathbf{b}_\perp) = \int \frac{d^2\mathbf{\Delta}_\perp}{(2\pi)^2} e^{i\mathbf{\Delta}_\perp \cdot \mathbf{b}_\perp} \tilde{H}(x, 0, -\mathbf{\Delta}_\perp^2).$$

Due to a Galilean subgroup of transverse boosts in the infinite momentum frame there are no relativistic corrections to Eq. (5). Furthermore, these impact parameter dependent parton distributions have a probabilistic interpretation very similar (and with the same limitations) as the usual PDFs[2].

So far, only few experiments exist that help constrain GPDs and therefore it is important to utilize theretical constraints when parameterizing these functions. One such constraint arises directly from the fact that the reference point for impact parameter dependent PDFs is the $\perp$ center of momentum. In the limit of $x \to 1$ the active quark becomes the center of momentum and therefore $\mathbf{b}_\perp$ can never be large for large x. For impact parameter dependent parton distributions this implies that their $\perp$ width should go to zero for $x \to 1$. For decreasing x the $\perp$ width is expected to gradually increase. Although the width in the valence region should still be relatively compact, its size should increase further once x is small enough for the pion cloud to contribute [3]. In momentum space, this implies that the t-dependence of GPDs should decrease with increasing x. This prediction is consistent with recent lattice results, which showed that higher moments of GPDs have less t dependence than lower moments [4]

3. Transversely Polarized Target

For a transversely polarized target, the impact parameter dependent parton distributions are no longer axially symmetric. The deviations from axial symmetry are



Fig. 1. Distribution of the j^+ density for u and d quarks in the $\perp$ plane ($x_{Bj} = 0.3$ is fixed) for a nucleon that is polarized in the x direction in the model from Ref. For other values of x the distortion looks similar. More recent fits yield similar results.

described by $E(x, 0, t)$. For example, the unpolarized quark distribution $q_X(x, \mathbf{b}_\perp)$ for a target that is polarized in the $+\hat{x}$ direction reads [5]

$$q_X(x,\mathbf{b}_\perp) = q(x,\mathbf{b}_\perp) - \frac{1}{2M}\frac{\partial}{\partial b_y}\int\frac{d^2\mathbf{\Delta}_\perp}{(2\pi)^2}E(x,0,-\mathbf{\Delta}_\perp^2)e^{-i\mathbf{b}_\perp\cdot\mathbf{\Delta}_\perp}. \tag{6}$$

Here $q(x,\mathbf{b}_\perp)$ is the impact parameter dependent PDF in the unpolarized case (5). The origin of this distortion is that the virtual photon in DIS couples more strongly to quarks that move towards it than quarks that move away from it (hence the γ^+ in the quark correlation function relevant for DIS). If the orbital motion of the quarks and the spin of the target are correlated then quarks are more likely to move towards the virtual photon on one side of the target than the other and therefore the distribution of quarks in impact parameter space appears deformed towards one side. The details of this deformation for each quark flavor are described by $E_q(x, 0, t)$, which is not known yet. However, sign and overall scale can be estimated by considering the mean displacement of flavor q ($\perp$ flavor dipole moment)

$$d_y^q \equiv \int dx \int d^2\mathbf{b}_\perp q(x,\mathbf{b}_\perp) b_y = \frac{1}{2M}\int dx E_q(x,0,0) = \frac{\kappa_q^p}{2M}. \tag{7}$$

The $\kappa_q = \mathcal{O}(1-2)$ are the anomalous magnetic moment contribution from each quark flavor to the anomalous magnetic moment of the nucleon, i.e. $F_2(0) = \frac{2}{3}\kappa_u - \frac{1}{3}\kappa_d - \frac{1}{3}\kappa_s ....$ This yields $\left|d_y^q\right| = \mathcal{O}(0.2fm)$, where u and d quarks have opposite signs. This is a sizeable effect as is illustrated in Fig. (1).

This deformation provides a very physical source for single-spin asymmetries (SSA) in semi-inclusive DIS. For an (on average) attractive final state interaction, the position space deformation into the $+\hat{y}$ direction translates into a momentum space asymmetry for the ejected quark that prefers the $-\hat{y}$ direction and *vice versa* (Fig. 2) Since the sign of the position space distortion is governed by the sign of the anomalous magnetic moment contribution $\kappa_{q/P}$ from each quark flavor, this implies that the sign of the SSA is correlated to the sign of $\kappa_{q/P}$. Following the Trento convention [7], this yields a negative Sivers function $f_{1T}^{\perp u}$ for u quarks in the proton, while $f_{1T}^{\perp d} > 0$. For the neutron those signs are reversed. These predictions are consistent with recent HERMES data [8].



Fig. 2. The transverse distortion of the parton cloud for a proton that is polarized into the plane, in combination with attractive FSI, gives rise to a Sivers effect for u (d) quarks with a $\perp$ momentum that is on the average up (down).

4. Chirally Odd GPDs

The distribution of transversely polarized quarks in impact parameter space is descibed by the Fourier transform of chirally odd GPDs [9]. For example, for an unpolarized target the distribution of quarks with transversity s^i reads

$$q_i(x,\mathbf{b}_\perp) = -\frac{s^i\varepsilon^{ij}}{2M}\frac{\partial}{2b_j}\int\frac{d^2\mathbf{\Delta}_\perp}{(2\pi)^2}\left[2\tilde{H}_T(x,0,-\mathbf{\Delta}_\perp^2)+E_T(x,0,-\mathbf{\Delta}_\perp^2)\right]e^{-i\mathbf{b}_\perp\cdot\mathbf{\Delta}_\perp}. \quad (8)$$

A consequence of this result is an analog of Ji's angular momentum sum rule, where the nucleon spin is replaced by the quark spin: the angular momentum J_q^i carried by quarks with transversity s^j in an unpolarized target reads [10]

$$\left\langle J_q^i(s^j)\right\rangle = \frac{\delta^{ij}}{4}\int dx\, x\left[H_T(x,0,0)+2\tilde{H}_T^q(x,0,0)+E_T^q(x,0,0)\right]. \quad (9)$$

The relation between Eq. (9) and Ji's sum rule [11] is similar to the relation between the Sivers function and the Boer-Mulders function $h_1^{\perp q}$ [12] in that the nucleon spin is replaced by the quark spin. This observation suggests extending the physical mechanism for the Sivers effect (Fig. 2) to the Boer-Mulders effect, yielding opposite signs for $2\tilde{H}_T^q(x,0,0)+E_T^q(x,0,0)$ and $h_1^{\perp q}$.

Acknowledgments

This work has been supported by the DOE (grant number DE-FG03-95ER40965).

References

1. M. Burkardt, Phys. Rev. **D62**, 071503 (2000); Erratum-ibid. **D66**, 119903 (2002).
2. M. Burkardt, hep-ph/0105324; P.V. Pobilitsa, Phys. Rev. **D70**, 034004 (2004).
3. M. Strikman and C. Weiss, Phys. Rev. **D69**, 054012 (2004).
4. Ph. Hägler et al., Phys. Rev. Lett. **93**, 112001 (2004).
5. M. Burkardt, Int. J. Mod. Phys. **A18**, 173 (2003).
6. X.Ji, Phys. Rev. Lett. **78**, 610 (2003).
7. A.Bacchetta et al., Phys. Rev. **D70**, 117504 (2004).
8. HERMES collaboration, Phys. Rev. Lett. **94**, 012002 (2005).
9. M. Diehl and Ph. Hägler, hep-ph/0504175.
10. M. Burkardt, hep-ph/0505189.
11. X. Ji, Phys. Rev. Lett. **78**, 610 (1997).
12. D. Boer and P.J. Mulders, Phys. Rev. **D57**, 5780 (1998).